

Synagis

CLIENT CARE PLAN

Client: \*Pathways , CarePlan

Client ID#:

1165425300

Admission Date:

12-06-2006

42 Medication Regimen

Potential Problem

Individual

Admission Rating	K	B	S
Desired Outcome	K	B	S

Cat.	Target	Care Description Note
TP	Medication Administration	Other Prescribed Medications IM RN to administer synagis IM to anterolateral aspect of thigh. (Children should NOT be given an injection in the gluteal muscle because of risk to the sciatic nerve)
TP	Medication Set-Up	Subcutaneous / Intramuscular Medications / Syringe Prefills RN to weigh baby. Synagis dosage is calculated as follows: Wt.(in KG) x 15mg/kg divided by 100mg/ml. Result is mg dosage to be given. Call pharmacy providing drug to double check calculations prior to administration of dosage. Injection volume >1ml should be given in divided doses. Synagis should be reconstituted with sterile water (per package instructions) and gently swirled for 30 sec to avoid foaming. DO NOT SHAKE. Let stand for 20 min until solution clarifies. Synagis should be administered within 5-6 hrs of reconstitution.
CM	Medication Set-Up	Other Order/Re-order Medications RN to coordinate delivery of monthly Synagis medication delivery to pt home prior to monthly injection
S	Medication Administration	Other RN to observe baby for 30min after injection to observe for s/sx of allergic or anaphylactic reactions. (Emergency drug kits are provided on first dose patients).
TGC	Medication Action/Side Effects	Other Synagis
TGC	Signs/Symptoms-Physical	When To Notify Providers Teach S/S anaphylaxis and/or S/S RSV to report

Employee Signature \_\_\_\_\_ Date \_\_\_\_\_